



## **Integrated micro/nanofibrous PLGA-Collagen scaffolds for bladder tissue regeneration**

**Ajalloueian, Fatemeh; Hilborn, Jons; Fossum, Magdalena; Chronakis, Ioannis S.**

*Published in:*  
Book of Abstracts. DTU's Sustain Conference 2015

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Ajalloueian, F., Hilborn, J., Fossum, M., & Chronakis, I. S. (2015). Integrated micro/nanofibrous PLGA-Collagen scaffolds for bladder tissue regeneration. In *Book of Abstracts. DTU's Sustain Conference 2015* [Q-13] Technical University of Denmark.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Integrated micro/nanofibrous PLGA-Collagen scaffolds for bladder tissue regeneration**

Fatemeh Ajalloueian<sup>1</sup>, Jons Hilborn<sup>2</sup>, Magdalena Fossum<sup>3</sup>, Ioannis S Chronakis<sup>1</sup>,

1: DTU Food, 2:Uppsala University, 3:Karolinska Institutet

\*Corresponding author email: [ioach@food.dtu.dk](mailto:ioach@food.dtu.dk)

Fabrication of an ideal scaffold mimicking the native extracellular matrix components, structure and function has remained a challenge in tissue engineering. The basic structural element in ECM is collagen type I with a fibrillar nano-scale structure. Collagen nanofibers can be fabricated using electrospinning or plastic compression (PC) of collagen hydrogel. PC is a simple and fast method, which has appeared better than electrospinning, since fluoroalcoholic solvents used in collagen electrospinning denature collagen to gelatin. The collagen layer obtained from PC contains small inter-fiber pores limiting cell infiltration and is mechanically weak. Performing collagen PC onto another polymeric substrate leads to a two-layered construct with improved mechanical properties. Here, we present the design of a hybrid, electrospun poly(lactic-co-glycolide) (PLGA) - plastically compressed (PC) collagen scaffold that could allow bladder mucosa expansion. Optimisation of electrospinning was performed in order to obtain increased pore sizes and porosity to consolidate the construct and to support neovascularisation and tissue ingrowth. The PLGA support scaffold was placed between two collagen gels, and the minced tissue was distributed either on top or both on top and inside the construct prior to PC; this was then cultured for up to four weeks. Morphology, histology and SEM demonstrated that the construct maintained its integrity throughout cell culture. Cells from minced tissue migrated, expanded and re-organised to a confluent cell layer on the top of the construct after two weeks and formed a multilayered urothelium after four weeks. Cell morphology and phenotypewas typical for urothelial mucosa during tissue culture.